

Equity-Oriented Treatment Agreements for Opiates or Controlled Drugs

Often, patients who are prescribed long term opiates or controlled drugs are asked to sign agreements or contracts to provide them with some parameters related to the ongoing use of these medications. Often, these agreements can convey a pejorative, patronizing or stigmatizing tone – even when unintended. To address this issue, clinicians at a primary care clinic recent rewrote their agreement template to be more equity oriented. They were able to institute these revisions immediately, and they report that both clinicians and patients are more comfortable with the revised version of the agreement. In this Tool, you will see the "before and after" versions of the agreement. In the marked up version, you can see the revisions the clinicians made to the agreement to reflect a more respectful and accepting tone. If you'd like to use this template for your own practice/organization, a clean copy of the revised version follows.

Sample Controlled Drug or Substance Agreement:

Introduction:

This is an agreement to help you use medication as safely as possible. This is for patients for whom we are planning to prescribe an opiate (also known as

a controlled drug). These medications have extra legal restrictions and have potentially harmful effects of use

some extra risks, including addiction. We ask all of our patients using one people who may need people or more of these medications, or planning to use these medications on a

regular basis, - to sign this agreement.

or may not work well for you

This medication may not be right for you. Everyone has a unique reaction

to the medication and the medications help people in unique ways. This

prescription will be started as a trial: If you don't notice much improvement in That is why it's useful to start this perscription on a trial basis:

a reasonable period of time we will help you gradually stop the medication.

using

If you transfer to our care, and are if you already taking controlled medications it may be helpful for us to work with you to adjust your medications for the following reasons: that don't seem to be helping you much; if you are taking combinations of harmful or

medications that are considered dangerous; or if you are taking higher than

currently recommended doses of a controlled drug. we will need to adjust help you adjust your medications

your medications. We will make medication adjustments gradually and

cautiously, to minimize any side effects you may feel as a result of changing your medications.

Very helpful phrasing and great way of explaining the purpose of the agreement.

If you look at the latest DSM-V terminology, you'll see that they reserve the term addiction only for gambling addiction.

Addiction is not used in relation to any substance use now. Hence this edit.

Removing 'ownership' and positioning people only as 'patients'

I understand and agree to the following:

		be harmful, for example, it can			
1		Receiving controlled drugs from multiple sources can result in overdose or harmful side effects of an accidental			
		dangerous drug combinations. For these reasons, it is important to chooget your medications from	se a		
		single primary care provider. It is also safer to use only one pharmacy.			
		The provider prescribing for you will be or	her/		
		his/their replacement. Pharmacy			
2	2.	Taking controlled medications in larger amounts or more frequently than			
		harms such as accidental overdose, or other harmful impacts on your health. prescribed is dangerous and can result in overdose. Take only the dose			
		prescribed and at time intervals prescribed. If you think you need your do	ose		
		or timing changed, discuss this with your provider.			
	В	Because everyone has a unique reaction to			
3		Gontrolled medications are prescribed carefully for an individual, and it especially for you individually			
		they especially for you individually dangerous to use these medications without consulting your provider. Described to the specifically for you specifically for you for these reason take controlled medications that are not prescribed for you. Do not	s, do		
	,	your medications to anyone else, including family members. Make sure y	our		
		medications are kept in a safe place where no-one, especially children or	r		
	aı	adolescents, can take the medication without your knowledge. It is also in potentially harmful to people's health to sell or give these medications away.	llega		
2	1.	Other people may try to steal these medications, therefore, These medications can be the target of thieves. It is important to store y	our		
		medications in a secure location.			
5	5.	Some over-the-counter medications including codeine containing			
		medications; cold medicines, sleeping pills and alcohol or recreational de	rugs		
		can have dangerous interactions with many controlled drugs. Check with	n		
		your pharmacist or provider before using over the counter medications a	and		
		be open about your use of alcohol and street drugs so your provider can	give		
		you accurate advice.			
	_	under very specific conditions and for very specific reasons			

6. Controlled drugs are prescribed very cautiously. If your prescription is lost or (for whatever reason, including if you have stolen or runs out early because you used more medication than prescribed), be able to your provider will likely not prescribe extra medications, in part because

prescribers' practices are carefully monitored. For this reason, it is helpful to keep in touch with your provider if you need to adjust your medications to ensure you are benefitting as much as possible from these medications.

- 7. Routine, random, urine drug screening is a standard precaution to ensure you are using your medication, and not using other drugs to monitor if the medications are helping you, and to check to see if there are other drugs that you may have used which could be dangerous in combination with this medication. Your provider is expected, as a standard for the licence to prescribe controlled drugs, to ask you to undergo urine drug screening. We greatly appreciate your willingness to do this on occasion.
- 8. It may be necessary for safe use of this medication to prescribe small amounts frequently, up to as often as daily witnessed medication use. The amount of medication you receive at one time and frequency of visits will be in partnership with you and your provider because these drugs are so carefully controlled, decided by your NP and you, but your provider is responsible for the final decision on frequency and amount of medication.
- 9. In order for the different members of the health care team to provide you with the best care possible, it is very helpful for them to have access to the records regarding the types of medication you take. I consent to open communication between my provider and any other health professionals involved in my pain management, such as pharmacists, other doctors, or emergency departments. This includes reviewing information available from PharmaNet.
- 10. If you are unable to use this medication safely and as prescribed, your

 This will be done in collaboration with you, and provider is obliged to discontinue the medication. Whenever possible this will be done by slowly tapering you off the medication to avoid withdrawal symptoms.

To signal that you are giving them a sense of control over the situation.

Patient's signature:	Clinician's signature:	
_	-	
Date:		

How to cite this document

EQUIP Health Care. (2017). Equity-Oriented Treatment Agreements for Opiates or Controlled Drugs. Vancouver, BC. Retrieved from www.equiphealthcare.ca.

Version | December 2017

References

Adapted From

https://fhs.mcmaster.ca/npc/opioid/cgop_b_app_b05.html

Canadian Guideline

http://www.cfp.ca/content/57/11/1257.full.pdf

FMPE (McMaster 2011) APPENDIX 3.

Sample Opioid Treatment Agreement

EQUIP Health Care & the Health Equity Toolkit are funded by CIHR

To learn more about EQUIP Health Care, please visit www.equiphealthcare.ca

To provide feedback about this tool, please visit https://www.surveymonkey.com/r/VRBF7WJ



